

PRAC antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14530**Specification**

PRAC antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q96KF2
Other Accession	NM_032391 , NP_115767
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	6kDa KDa

PRAC antibody - N-terminal region - Additional Information**Gene ID** 84366**Alias Symbol** **C17orf92, MGC32520****Other Names**

Small nuclear protein PRAC1, Prostate cancer susceptibility candidate protein 1, Prostate, rectum and colon expressed gene protein, PRAC1, C17orf92, PRAC

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PRAC antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

PRAC antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PRAC antibody - N-terminal region - Protein Information**Name** PRAC1**Synonyms** C17orf92, PRAC**Cellular Location**

Nucleus.

Tissue Location


Highly expressed in prostate, rectum, and distal colon, and weakly expressed in bladder. Expressed in prostate cancer cell lines.

PRAC antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PRAC antibody - N-terminal region - Images



WB Suggested Anti-PRAC Antibody Titration: 1.0 µg/ml
Positive Control: Fetal Liver

PRAC antibody - N-terminal region - References

Liu X.F., et al. Prostate 47:125-131(2001).
Zody M.C., et al. Nature 440:1045-1049(2006).
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.