

RIOK1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14538**Specification**

RIOK1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9H2L9
Other Accession	NM_153005 , NP_694550
Reactivity	Human, Horse
Predicted	Human, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38kDa KDa

RIOK1 antibody - C-terminal region - Additional Information

Alias Symbol	AD034, FLJ30006, MGC12903, bA288G3.1, RRP10
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-RIOK1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


RIOK1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

RIOK1 antibody - C-terminal region - Protein Information**RIOK1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RIOK1 antibody - C-terminal region - Images



WB Suggested Anti-R1OK1 Antibody Titration: 1.0 µg/ml
Positive Control: PANC1 Whole Cell