

**CD99L2 antibody - middle region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI14540****Specification**

---

**CD99L2 antibody - middle region - Product Information**

|                   |                                                                |
|-------------------|----------------------------------------------------------------|
| Application       | WB                                                             |
| Primary Accession | <a href="#">Q8TCZ2</a>                                         |
| Other Accession   | <a href="#">NM_134446</a> , <a href="#">NP_604395</a>          |
| Reactivity        | Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog |
| Predicted Host    | Human, Mouse, Rabbit, Pig, Horse, Dog                          |
| Clonality         | Rabbit                                                         |
| Calculated MW     | Polyclonal<br>23kDa kDa                                        |

**CD99L2 antibody - middle region - Additional Information****Gene ID** 83692**Alias Symbol** CD99B, DKFZp761H2024, MIC2L1**Other Names**

CD99 antigen-like protein 2, MIC2-like protein 1, CD99, CD99L2, MIC2L1

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-CD99L2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

CD99L2 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

**CD99L2 antibody - middle region - Protein Information****Name** CD99L2**Synonyms** MIC2L1**Function**

Plays a role in a late step of leukocyte extravasation helping cells to overcome the endothelial basement membrane. Acts at the same site as, but independently of, PECAM1 (By similarity). Homophilic adhesion molecule, but these interactions may not be required for cell aggregation (By similarity).

**Cellular Location**

Cell membrane; Single-pass type I membrane protein; Extracellular side. Cell junction. Secreted


**Tissue Location**

Detected in cerebrospinal fluid (at protein level) (PubMed:25326458). Expressed in many tissues, with low expression in thymus.

**CD99L2 antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CD99L2 antibody - middle region - Images**

WB Suggested Anti-CD99L2 Antibody Titration: 1.0 µg/ml  
Positive Control: PANC1 Whole Cell