

ACOT9 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14575**Specification**

ACOT9 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O96EA2
Other Accession	NM_001033583 , NP_001028755
Reactivity	Human, Rat, Horse, Bovine, Guinea Pig
Predicted	Human, Pig, Horse, Bovine, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	23kDa KDa

ACOT9 antibody - C-terminal region - Additional Information

Alias Symbol **ACATE2, CGI-16, MT-ACT48, MTACT48**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ACOT9 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


ACOT9 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ACOT9 antibody - C-terminal region - Protein Information**ACOT9 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ACOT9 antibody - C-terminal region - Images



WB Suggested Anti-ACOT9 Antibody Titration: 1.0 µg/ml
Positive Control: MDA-MB-435S Whole Cell