

Rab5a antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14626**Specification**

Rab5a antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9COD1
Other Accession	NM_025887 , NP_080163
Reactivity	Human, Mouse, Rat, Rabbit, Sheep, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Pig, Chicken, Sheep, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24kDa KDa

Rab5a antibody - C-terminal region - Additional Information**Gene ID** 271457

Alias Symbol	2410015H04Rik, AI663973, AU021172
Other Names	
Ras-related protein Rab-5A, Rab5a, nnyRab5a	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Rab5a antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Rab5a antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Rab5a antibody - C-terminal region - Protein Information**Name** Rab5a {ECO:0000312|MGI:MGI:105926}**Synonyms** nnyRab5a**Function**

The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. RAB5A is required for the fusion of plasma membranes and early endosomes. Contributes

to the regulation of filopodia extension. Required for the exosomal release of SDCBP, CD63, PDCD6IP and syndecan (By similarity). Regulates maturation of apoptotic cell-containing phagosomes, probably downstream of DYN2 and PIK3C3 (PubMed:18425118).

Cellular Location


Cell membrane {ECO:0000250|UniProtKB:P20339}; Lipid-anchor {ECO:0000250|UniProtKB:P20339}; Cytoplasmic side {ECO:0000250|UniProtKB:P18066}. Early endosome membrane {ECO:0000250|UniProtKB:P20339}; Lipid-anchor {ECO:0000250|UniProtKB:P20339}. Melanosome. Cytoplasmic vesicle {ECO:0000250|UniProtKB:P20339}. Cell projection, ruffle {ECO:0000250|UniProtKB:P18066}. Membrane {ECO:0000250|UniProtKB:P20339}. Cytoplasm, cytosol {ECO:0000250|UniProtKB:P20339}. Cytoplasmic vesicle, phagosome membrane. Endosome membrane {ECO:0000250|UniProtKB:P20339}. Note=Enriched in stage I melanosomes Alternates between membrane-bound and cytosolic forms {ECO:0000250|UniProtKB:P20339}

Rab5a antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Rab5a antibody - C-terminal region - Images



WB Suggested Anti-Rab5a Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Small Intestine

Rab5a antibody - C-terminal region - References

Kuroda T.S.,et al.J. Biol. Chem. 277:9212-9218(2002).
Itoh T.,et al.Genes Cells 11:1023-1037(2006).
Carninci P.,et al.Science 309:1559-1563(2005).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Lubec G.,et al.Submitted (APR-2007) to UniProtKB.