

DTNA antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14676**Specification**

DTNA antibody - C-terminal region - Product Information

Application	WB
Primary Accession	B4DIRO
Other Accession	NM_001198942 , NP_001185871
Reactivity	Human, Mouse, Rat, Horse, Yeast, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	49kDa kDa

DTNA antibody - C-terminal region - Additional Information

Alias Symbol **D18S892E, DRP3, DTN, FLJ96209, LVNC1**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-DTNA antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


DTNA antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DTNA antibody - C-terminal region - Protein Information**DTNA antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DTNA antibody - C-terminal region - Images



WB Suggested Anti-DTNA Antibody Titration: 1.0 µg/ml
Positive Control: Hela Whole Cell