

PAF1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14679**Specification**

PAF1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8N7H5
Other Accession	NM_019088 , NP_061961
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	58kDa kDa

PAF1 antibody - N-terminal region - Additional Information**Gene ID** 54623**Alias Symbol** **F23149_1, FLJ11123, PD2****Other Names**

RNA polymerase II-associated factor 1 homolog, hPAF1, Pancreatic differentiation protein 2, PAF1, PD2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PAF1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

PAF1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PAF1 antibody - N-terminal region - Protein Information**Name** PAF1**Synonyms** PD2**Function**

Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser- 5'-phosphorylated forms and is involved in

transcriptional elongation, acting both independently and synergistically with TCEA1 and in cooperation with the DSIF complex and HTATSF1. PAF1C is required for transcription of Hox and Wnt target genes. PAF1C is involved in hematopoiesis and stimulates transcriptional activity of KMT2A/MLL1; it promotes leukemogenesis through association with KMT2A/MLL1-rearranged oncoproteins, such as KMT2A/MLL1-MLLT3/AF9 and KMT2A/MLL1-MLLT1/ENL. PAF1C is involved in histone modifications such as ubiquitination of histone H2B and methylation on histone H3 'Lys-4' (H3K4me3). PAF1C recruits the RNF20/40 E3 ubiquitin-protein ligase complex and the E2 enzyme UBE2A or UBE2B to chromatin which mediate monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1); UB2A/B-mediated H2B ubiquitination is proposed to be coupled to transcription. PAF1C is involved in mRNA 3' end formation probably through association with cleavage and poly(A) factors. In case of infection by influenza A strain H3N2, PAF1C associates with viral NS1 protein, thereby regulating gene transcription. Connects PAF1C with the RNF20/40 E3 ubiquitin-protein ligase complex. Involved in polyadenylation of mRNA precursors. Has oncogenic activity in vivo and in vitro.

Cellular Location


Nucleus. Note=Punctuate distribution throughout the nucleus except in nucleoli and the perinuclear chromatin

PAF1 antibody - N-terminal region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PAF1 antibody - N-terminal region - Images



WB Suggested Anti-PAF1 Antibody Titration: 1.0 µg/ml
Positive Control: HepG2 Whole Cell




Host:Rabbit

Target Name:PAF1

Sample Tissue:Human 721_B

Antibody Dilution: 1.0µg/ml PAF1 is supported by BioGPS gene expression data to be expressed in 721_B




Host:Rabbit

Target Name:PAF1

Sample Tissue:Human HeLa

Antibody Dilution: 1.0µg/ml PAF1 is supported by BioGPS gene expression data to be expressed in HeLa



Host:Rabbit

Target Name:PAF1

Sample Tissue:Human Fetal Heart

Antibody Dilution: 1.0µg/ml

PAF1 antibody - N-terminal region - References

Moniaux N.,et al.Oncogene 25:3247-3257(2006).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Grimwood J.,et al.Nature 428:529-535(2004).

Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Zhu B.,et al.Genes Dev. 19:1668-1673(2005).