

DOK7 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14721**Specification**

DOK7 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q18PE1-2
Other Accession	NM_173660 , NP_775931
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40kDa KDa

DOK7 antibody - C-terminal region - Additional Information

Alias Symbol	C4orf25, CMS1B, FLJ33718, FLJ39137, FLJ90556
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-DOK7 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


DOK7 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DOK7 antibody - C-terminal region - Protein Information**DOK7 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DOK7 antibody - C-terminal region - Images



WB Suggested Anti-DOK7 Antibody Titration: 1.0 µg/ml
Positive Control: Jurkat Whole Cell