

Tacc2 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14738**Specification**

Tacc2 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9JIG0-4
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	65kDa KDa

Tacc2 Antibody - C-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Tacc2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


Tacc2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Tacc2 Antibody - C-terminal region - Protein Information**Tacc2 Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Tacc2 Antibody - C-terminal region - Images



168 kDa__
144 kDa__
90 kDa__
65 kDa__
40 kDa__

Host: Rabbit
Target Name: Tacc2
Sample Tissue: Mouse Muscle lysates
Antibody Dilution: 1.0µg/ml