

TAGAP antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14783**Specification**

TAGAP antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O8N103-2
Other Accession	NM_152133 , NP_687034
Reactivity	Human, Guinea Pig
Predicted	Human, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60kDa KDa

TAGAP antibody - C-terminal region - Additional Information

Alias Symbol	FKSG15, FLJ32631, FLJ39771, IDDM21, MGC133247, MGC27381, TAGAP1, ARHGAP47
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-TAGAP antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


TAGAP antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TAGAP antibody - C-terminal region - Protein Information**TAGAP antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TAGAP antibody - C-terminal region - Images



WB Suggested Anti-TAGAP Antibody Titration: 1.0 µg/ml
Positive Control: MCF7 Whole Cell