

ADAT3 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14824**Specification**

ADAT3 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q96EY9
Other Accession	NM_138422 , NP_612431
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39kDa KDa

ADAT3 antibody - C-terminal region - Additional Information**Gene ID** 113179**Alias Symbol** FWP005, MST121, MSTP121, S863-5, TAD3**Other Names**

Probable inactive tRNA-specific adenosine deaminase-like protein 3, tRNA-specific adenosine-34 deaminase subunit ADAT3, ADAT3, TAD3

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ADAT3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ADAT3 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


ADAT3 antibody - C-terminal region - Protein Information**Name** ADAT3**Synonyms** TAD3**ADAT3 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ADAT3 antibody - C-terminal region - Images



WB Suggested Anti-ADAT3 Antibody Titration: 1.0 µg/ml
Positive Control: Jurkat Whole Cell

ADAT3 antibody - C-terminal region - References

Gauci S., et al. Anal. Chem. 81:4493-4501(2009).
Burkard T.R., et al. BMC Syst. Biol. 5:17-17(2011).
Sjoeblom T., et al. Science 314:268-274(2006).
Alazami A.M., et al. J. Med. Genet. 50:425-430(2013).