

SLAMF1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14855**Specification**

SLAMF1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	B2RAL4
Other Accession	NM_003037 , NP_003028
Reactivity	Human, Pig
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37kDa KDa

SLAMF1 antibody - C-terminal region - Additional Information

Alias Symbol **CD150, CDw150, SLAM**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SLAMF1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

SLAMF1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SLAMF1 antibody - C-terminal region - Protein Information**SLAMF1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SLAMF1 antibody - C-terminal region - Images




90 kDa
65 kDa
40 kDa
29 kDa
22 kDa

WB Suggested Anti-SLAMF1 Antibody Titration: 1.0 μ g/ml

Positive Control: 721_B Whole Cell SLAMF1 is supported by BioGPS gene expression data to be expressed in 721_B