

WBSCR22 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14860**Specification**

WBSCR22 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	C9K060
Other Accession	NM_001202560 , NP_001189489
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32kDa KDa

WBSCR22 antibody - C-terminal region - Additional Information

Alias Symbol	HASJ4442, HUSSY-3, MGC19709, MGC2022, MGC5140, PP3381, WBMT
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-WBSCR22 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


WBSCR22 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

WBSCR22 antibody - C-terminal region - Protein Information**WBSCR22 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

WBSCR22 antibody - C-terminal region - Images



WB Suggested Anti-WBSCR22 Antibody Titration: 1.0 µg/ml

Positive Control: RPMI-8226 Whole Cell
WBSCR22 is supported by BioGPS gene expression data to be expressed in RPMI 8226