

ZFP62 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14921**Specification**

ZFP62 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8NB50
Other Accession	NM_152283 , NP_689496
Reactivity	Human, Rabbit, Bovine
Predicted	Human, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	103kDa KDa

ZFP62 antibody - C-terminal region - Additional Information**Gene ID** 643836**Alias Symbol** DKFZp667F2013, FLJ11344, FLJ34057, FLJ34231, FLJ58781, FLJ59694, MGC176438, ZET, ZNF755**Other Names**

Zinc finger protein 62 homolog, Zfp-62, ZFP62

Target/Specificity

Recognizes both isoforms of human ZFP62.

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZFP62 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZFP62 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZFP62 antibody - C-terminal region - Protein Information**Name** ZFP62**Function**

May play a role in differentiating skeletal muscle.

Cellular Location


Nucleus.

ZFP62 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZFP62 antibody - C-terminal region - Images



WB Suggested Anti-ZFP62 Antibody Titration: 1.0 µg/ml
Positive Control: RPMI-8226 Whole Cell

ZFP62 antibody - C-terminal region - References

Ota T., et al. Nat. Genet. 36:40-45(2004).
Schmutz J., et al. Nature 431:268-274(2004).
Bechtel S., et al. BMC Genomics 8:399-399(2007).