

SYNC antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14991**Specification**

SYNC antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9H7C4-2
Other Accession	NM_001161708 , NP_001155180
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52kDa KDa

SYNC antibody - C-terminal region - Additional Information

Alias Symbol	MGC149625, MGC149626, SYNC1, SYNCOILIN
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SYNC antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

SYNC antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SYNC antibody - C-terminal region - Protein Information**SYNC antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SYNC antibody - C-terminal region - Images

WB Suggested Anti-SYNC Antibody Titration: 1.0 μ g/ml
Positive Control: U937 Whole Cell