

CDC73 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # Al14995

Specification

CDC73 antibody - C-terminal region - Product Information

Application WB
Primary Accession Q6P1|9

Other Accession NM 024529, NP 078805

Reactivity Human, Mouse, Rat, Rabbit, Horse, Bovine,

Dog

Predicted Human, Mouse, Rat, Chicken, Horse,

Bovine, Dog

Host Rabbit
Clonality Polyclonal
Calculated MW 58kDa KDa

CDC73 antibody - C-terminal region - Additional Information

Gene ID 79577

Alias Symbol

Clorf28, FLJ23316, HPTJT, HRPT2, HYX

Other Names

Parafibromin, Cell division cycle protein 73 homolog, Hyperparathyroidism 2 protein, CDC73, C1orf28, HRPT2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CDC73 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CDC73 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CDC73 antibody - C-terminal region - Protein Information

Name CDC73

Synonyms Clorf28, HRPT2

Function

Tumor suppressor probably involved in transcriptional and post-transcriptional control pathways. May be involved in cell cycle progression through the regulation of cyclin D1/PRAD1 expression. Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and maintenance of embryonic stem



cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'-phosphorylated forms and is involved in transcriptional elongation, acting both independently and synergistically with TCEA1 and in cooperation with the DSIF complex and HTATSF1. PAF1C is required for transcription of Hox and Wnt target genes. PAF1C is involved in hematopoiesis and stimulates transcriptional activity of KMT2A/MLL1; it promotes leukemogenesis through association with KMT2A/MLL1-rearranged oncoproteins, such as KMT2A/MLL1-MLLT3/AF9 and KMT2A/MLL1-MLLT1/ENL. PAF1C is involved in histone modifications such as ubiquitination of histone H2B and methylation on histone H3 'Lys-4' (H3K4me3). PAF1C recruits the RNF20/40 E3 ubiquitin-protein ligase complex and the E2 enzyme UBE2A or UBE2B to chromatin which mediate monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1); UB2A/B-mediated H2B ubiquitination is proposed to be coupled to transcription. PAF1C is involved in mRNA 3' end formation probably through association with cleavage and poly(A) factors. In case of infection by influenza A strain H3N2, PAF1C associates with viral NS1 protein, thereby regulating gene transcription. Connects PAF1C with the cleavage and polyadenylation specificity factor (CPSF) complex and the cleavage stimulation factor (CSTF) complex, and with Wnt signaling. Involved in polyadenylation of mRNA precursors.

Cellular Location Nucleus

Tissue Location


Found in adrenal and parathyroid glands, kidney and heart.

CDC73 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CDC73 antibody - C-terminal region - Images



WB Suggested Anti-CDC73 Antibody Titration: 1.0 μg/ml

Positive Control: MDA-MB-435S Whole Cell

CDC73 antibody - C-terminal region - References





Tel: 858.875.1900 Fax: 858.875.1999

Sood R., et al. Genomics 73:211-222(2001). Ota T., et al. Nat. Genet. 36:40-45(2004). Gregory S.G., et al. Nature 441:315-321(2006). Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases. Bienvenut W.V., et al. Submitted (FEB-2008) to UniProtKB.