

INF2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15001**Specification**

INF2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q27J81
Other Accession	NM_022489 , NP_071934
Reactivity	Human, Mouse, Rat, Horse, Bovine, Dog
Predicted	Human, Mouse, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37kDa KDa

INF2 antibody - C-terminal region - Additional Information**Gene ID** 64423**Alias Symbol** C14orf151, C14orf173, DKFZp762A0214, FLJ22056, FSGS5, MGC13251, pp9484, CMTDIE**Other Names**

Inverted formin-2, HBEBP2-binding protein C, INF2, C14orf151, C14orf173

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-INF2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

INF2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

INF2 antibody - C-terminal region - Protein Information**Name** INF2**Synonyms** C14orf151, C14orf173**Function**

Severs actin filaments and accelerates their polymerization and depolymerization.

Cellular Location

Cytoplasm, perinuclear region


Tissue Location

Widely expressed. In the kidney, expression is apparent in podocytes and some tubule cells

INF2 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

INF2 antibody - C-terminal region - Images

WB Suggested Anti-INF2 Antibody Titration: 1.0 µg/ml

Positive Control: HepG2 Whole Cell INF2 is supported by BioGPS gene expression data to be expressed in HepG2