

PICALM antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15044**Specification**

PICALM antibody - C-terminal region - Product Information

Application	WB
Primary Accession	E9PN05
Other Accession	NM_001206947 , NP_001193876
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60kDa KDa

PICALM antibody - C-terminal region - Additional Information

Alias Symbol **CALM, CLTH, LAP**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PICALM antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


PICALM antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PICALM antibody - C-terminal region - Protein Information**PICALM antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PICALM antibody - C-terminal region - Images



WB Suggested Anti-PICALM Antibody Titration: 1.0 µg/ml
Positive Control: Jurkat Whole Cell