

**SNX1 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI15053****Specification****SNX1 antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">Q13596</a>
Other Accession	<a href="#">NM_001242933</a> , <a href="#">NP_001229862</a>
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63kDa KDa

**SNX1 antibody - N-terminal region - Additional Information****Gene ID 6642**

Alias Symbol	<a href="#">HsT17379</a> , <a href="#">MGC8664</a> , <a href="#">SNX1A</a> , <a href="#">Vps5</a> , <a href="#">VPS5</a>
Other Names	

Sorting nexin-1, SNX1

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-SNX1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

SNX1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**SNX1 antibody - N-terminal region - Protein Information****Name SNX1****Function**

Involved in several stages of intracellular trafficking. Interacts with membranes containing phosphatidylinositol 3-phosphate (PtdIns(3P)) or phosphatidylinositol 3,5-bisphosphate (PtdIns(3,5)P2) (PubMed:<a href="http://www.uniprot.org/citations/12198132" target="\_blank">12198132</a>). Acts in part as component of the retromer membrane-deforming SNX-BAR subcomplex. The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans-Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX-BAR subcomplex functions to deform the

donor membrane into a tubular profile called endosome-to-TGN transport carrier (ETC) (Probable). Can sense membrane curvature and has in vitro vesicle-to-membrane remodeling activity (PubMed:<a href="http://www.uniprot.org/citations/19816406" target="\_blank">19816406</a>, PubMed:<a href="http://www.uniprot.org/citations/23085988" target="\_blank">23085988</a>). Involved in retrograde endosome-to-TGN transport of lysosomal enzyme receptors (IGF2R, M6PR and SORT1) and Shigella dysenteria toxin stxB. Plays a role in targeting ligand-activated EGFR to the lysosomes for degradation after endocytosis from the cell surface and release from the Golgi (PubMed:<a href="http://www.uniprot.org/citations/12198132" target="\_blank">12198132</a>, PubMed:<a href="http://www.uniprot.org/citations/15498486" target="\_blank">15498486</a>, PubMed:<a href="http://www.uniprot.org/citations/17101778" target="\_blank">17101778</a>, PubMed:<a href="http://www.uniprot.org/citations/17550970" target="\_blank">17550970</a>, PubMed:<a href="http://www.uniprot.org/citations/18088323" target="\_blank">18088323</a>, PubMed:<a href="http://www.uniprot.org/citations/21040701" target="\_blank">21040701</a>). Involvement in retromer-independent endocytic trafficking of P2RY1 and lysosomal degradation of protease-activated receptor-1/F2R (PubMed:<a href="http://www.uniprot.org/citations/16407403" target="\_blank">16407403</a>, PubMed:<a href="http://www.uniprot.org/citations/20070609" target="\_blank">20070609</a>). Promotes KALRN- and RHOG-dependent but retromer-independent membrane remodeling such as lamellipodium formation; the function is dependent on GEF activity of KALRN (PubMed:<a href="http://www.uniprot.org/citations/20604901" target="\_blank">20604901</a>). Required for endocytosis of DRD5 upon agonist stimulation but not for basal receptor trafficking (PubMed:<a href="http://www.uniprot.org/citations/23152498" target="\_blank">23152498</a>).

### Cellular Location


Endosome membrane; Peripheral membrane protein; Cytoplasmic side. Golgi apparatus, trans-Golgi network membrane; Peripheral membrane protein; Cytoplasmic side. Early endosome membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, lamellipodium. Note=Enriched on tubular elements of the early endosome membrane. Binds preferentially to highly curved membranes enriched in phosphatidylinositol 3-phosphate (PtdIns(3P)) or phosphatidylinositol 3,5-bisphosphate (PtdIns(3,5)P2) (PubMed:15498486). Colocalized with SORT1 to tubular endosomal membrane structures called endosome-to-TGN transport carriers (ETCs) which are budding from early endosome vacuoles just before maturing into late endosome vacuoles (PubMed:18088323). Colocalizes with DNAJC13 and Shigella dysenteria toxin stxB on early endosomes (PubMed:19874558) Colocalized with F-actin at the leading edge of lamellipodia in a KALRN-dependent manner (PubMed:20604901).

### SNX1 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### SNX1 antibody - N-terminal region - Images



WB Suggested Anti-SNX1 Antibody Titration: 1.0 µg/ml

Positive Control: 721\_B Whole Cells SNX1 is strongly supported by BioGPS gene expression data to be expressed in Human 721\_B cells



Human , Mouse

#### **SNX1 antibody - N-terminal region - References**

- Kurten R.C.,et al.Science 272:1008-1010(1996).  
Haft C.R.,et al.Mol. Cell. Biol. 18:7278-7287(1998).  
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Zody M.C.,et al.Nature 440:671-675(2006).