

LTB4R2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15085**Specification**

LTB4R2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q5KU28
Other Accession	NM_001164692 , NP_001158164
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Bovine
Predicted	Human, Rabbit, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38kDa KDa

LTB4R2 antibody - C-terminal region - Additional Information

Alias Symbol **BLT2, BLTR2, JULF2, KPG_004, NOP9**
Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-LTB4R2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


LTB4R2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

LTB4R2 antibody - C-terminal region - Protein Information**LTB4R2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

LTB4R2 antibody - C-terminal region - Images



90 kDa
65 kDa
40 kDa
29 kDa
22 kDa

WB Suggested Anti-LTB₄R2 Antibody Titration: 1.0 µg/ml

Positive Control: 721_B Whole CellLTB₄R2 is supported by BioGPS gene expression data to be expressed in 721_B