

RBFA antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15160**Specification**

RBFA antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8N0V3
Other Accession	NM_024805 , NP_079081
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34kDa KDa

RBFA antibody - C-terminal region - Additional Information**Gene ID** 79863**Alias Symbol** **FLJ18281, FLJ18505, FLJ21172, FLJ42604, FLJ97160, HsT169****Other Names**

Putative ribosome-binding factor A, mitochondrial, RBFA, C18orf22

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-RBFA antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

RBFA antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

RBFA antibody - C-terminal region - Protein Information**Name** RBFA**Synonyms** C18orf22**Cellular Location**


Mitochondrion.

RBFA antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RBFA antibody - C-terminal region - Images



WB Suggested Anti-RBFA Antibody Titration: 1.0 μ g/ml

Positive Control: Hela Whole Cell

RBFA antibody - C-terminal region - References

Wang C.,et al.Submitted (OCT-2001) to the EMBL/GenBank/DDBJ databases.

Ota T.,et al.Nat. Genet. 36:40-45(2004).