

MON2 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15276**Specification**

MON2 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q6P148
Other Accession	NM_015026 , AAH65289
Reactivity	Human, Rat, Horse, Bovine, Dog
Predicted	Human, Rat, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	72kDa KDa

MON2 Antibody - C-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-MON2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


MON2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

MON2 Antibody - C-terminal region - Protein Information**MON2 Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MON2 Antibody - C-terminal region - Images



WB Suggested Anti-MON2 Antibody Titration: 1.0 µg/ml
Positive Control: Fetal Kidney