

LRRC10 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15399**Specification**

LRRC10 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q5BKY1
Other Accession	NM_201550 , NP_963844
Reactivity	Human, Rabbit, Bovine
Predicted	Human, Rabbit, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30kDa KDa

LRRC10 Antibody - C-terminal region - Additional Information**Gene ID** 376132**Alias Symbol** HRLRRP, LRRC10A, MGC125812**Other Names**

Leucine-rich repeat-containing protein 10, LRRC10

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-LRRC10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

LRRC10 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

LRRC10 Antibody - C-terminal region - Protein Information**Name** LRRC10**Function**

May play important roles in cardiac development and/or cardiac function.

Cellular Location


Nucleus.

LRRC10 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

LRRC10 Antibody - C-terminal region - Images



WB Suggested Anti-LRRC10 Antibody Titration: 1.0 µg/ml

Positive Control: PANC1 Whole Cell

LRRC10 Antibody - C-terminal region - References

Ota T., et al. Nat. Genet. 36:40-45(2004).