

ISOC1 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15434**Specification**

ISOC1 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O96CN7
Other Accession	NM_016048 , NP_057132
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22kDa KDa

ISOC1 Antibody - C-terminal region - Additional Information**Gene ID** 51015

Alias Symbol	CGI-111
Other Names	
Isochorismatase domain-containing protein 1, ISOC1	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ISOC1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ISOC1 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


ISOC1 Antibody - C-terminal region - Protein Information**Name** ISOC1**ISOC1 Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ISOC1 Antibody - C-terminal region - Images



WB Suggested Anti-ISOC1 Antibody Titration: 1.0 µg/ml
Positive Control: Fetal Heart

ISOC1 Antibody - C-terminal region - References

Burkard T.R., et al. BMC Syst. Biol. 5:17-17(2011).