

STK35 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15466**Specification**

STK35 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8TDR2
Other Accession	NM_080836 , NP_543026
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44kDa kDa

STK35 Antibody - C-terminal region - Additional Information**Gene ID** 140901**Alias Symbol** CLIK1, STK35L1**Other Names**

Serine/threonine-protein kinase 35, 2.7.11.1, CLP-36-interacting kinase 1, CLIK-1, PDLIM1-interacting kinase 1, Serine/threonine-protein kinase 35 L1, STK35, CLIK1, PDIK1, STK35L1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-STK35 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

STK35 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

STK35 Antibody - C-terminal region - Protein Information**Name** STK35**Synonyms** CLIK1, PDIK1, STK35L1**Cellular Location**

Nucleus. Nucleus, nucleolus. Cytoplasm. Note=When associated with PDLIM1, it is mostly found in cytoplasm, localized to actin stress fibers (PubMed:11973348). However, PubMed:19756140 detected STK35 only in the nucleus, and the presence of PDLIM1 had no influence on its location.


Tissue Location
Expressed in testis.

STK35 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STK35 Antibody - C-terminal region - Images



Host: Rabbit
Target Name: STK35
Sample Tissue: A549 Whole cell lysate
s
Antibody Dilution: 1.0µg/ml