

**CLEC10A Antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI15484**

**Specification**

**CLEC10A Antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">Q8IUN9</a>
Other Accession	<a href="#">NP_878910</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34kDa KDa

**CLEC10A Antibody - N-terminal region - Additional Information**

**Gene ID** 10462

**Alias Symbol**

**CLEC10A, CLECSF13, CLECSF14, HML,**

**Other Names**

C-type lectin domain family 10 member A, C-type lectin superfamily member 14, Macrophage lectin 2, CD301, CLEC10A, CLECSF13, CLECSF14, HML

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 &mu; l of distilled water. Final Anti-CLEC10A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

CLEC10A Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**CLEC10A Antibody - N-terminal region - Protein Information**

**Name** CLEC10A {ECO:0000303|PubMed:33724805}

**Function**

C-type lectin receptor involved in recognition of N- acetylgalactosamine (GalNAc)-terminated glycans by myeloid antigen presenting cells (APCs) (PubMed:<a href="http://www.uniprot.org/citations/15802303" target="\_blank">15802303</a>, PubMed:<a href="http://www.uniprot.org/citations/16998493" target="\_blank">16998493</a>, PubMed:<a href="http://www.uniprot.org/citations/17616966" target="\_blank">17616966</a>, PubMed:<a href="http://www.uniprot.org/citations/22213806" target="\_blank">22213806</a>, PubMed:<a href="http://www.uniprot.org/citations/33724805" target="\_blank">33724805</a>, PubMed:<a href="http://www.uniprot.org/citations/8598452" target="\_blank">8598452</a>). Binds in a

Ca(2+)-dependent manner to alpha- and beta-linked GalNAc residues on glycoprotein and glycolipid antigens, including alphaGalNAc- and Galbeta1->3GalNAc-O-Ser/Thr also known as Tn and T antigens, LacdiNAc epitope GalNAcbeta1->4GlcNAc and its derivative GalNAcbeta1->4-(Fucalpha1->3)GlcNAc, O-linked core 5 and 6 glycans, and GM2 and GD2 gangliosides (PubMed:<a href="http://www.uniprot.org/citations/15802303" target="\_blank">15802303</a>, PubMed:<a href="http://www.uniprot.org/citations/23507963" target="\_blank">23507963</a>). Acts as a signaling receptor at the interface of APC-T cell interactions. On immature dendritic cells, recognizes Tn antigen-carrying PTPRC/CD45 receptor on effector T cells and downregulates PTRPN/CD45 phosphatase activity with an impact on T cell activation threshold, cytokine production and proliferation. Modulates dendritic cell maturation toward a tolerogenic phenotype leading to generation of regulatory CD4-positive T cell subset with immune suppressive functions (PubMed:<a href="http://www.uniprot.org/citations/15802303" target="\_blank">15802303</a>, PubMed:<a href="http://www.uniprot.org/citations/16998493" target="\_blank">16998493</a>, PubMed:<a href="http://www.uniprot.org/citations/22213806" target="\_blank">22213806</a>). Acts as an endocytic pattern recognition receptor involved in antitumor immunity. During tumorigenesis, recognizes Tn antigens and its sialylated forms Neu5Ac-Tn and Neu5Gc-Tn expressed on tumor cell mucins. On immature dendritic cells, can internalize Tn-terminated immunogens and target them to endolysosomal compartment for MHC class I and II antigen presentation to CD8-positive and CD4-positive T cells, respectively (PubMed:<a href="http://www.uniprot.org/citations/15802303" target="\_blank">15802303</a>, PubMed:<a href="http://www.uniprot.org/citations/17616966" target="\_blank">17616966</a>, PubMed:<a href="http://www.uniprot.org/citations/17804752" target="\_blank">17804752</a>).

### **Cellular Location**

Cell membrane; Single-pass type II membrane protein. Early endosome membrane; Single-pass type II membrane protein Lysosome membrane; Single-pass type II membrane protein.

Note=Recycles between the plasma membrane and the endolysosomal compartment. Upon antigen binding, internalizes via endocytosis and then dissociates from antigen at acidic pH characteristic of endolysosomal vesicles

### **Tissue Location**


Expressed in myeloid antigen presenting cells in lymph nodes and skin (at protein level). Expressed in dermal dendritic cells (at protein level).

### **CLEC10A Antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **CLEC10A Antibody - N-terminal region - Images**



Host: Rabbit  
Target Name: CLEC10A  
Sample Tissue: HepG2 Whole cell lysate  
Antibody Dilution: 1.0 $\mu$ g/ml

#### **CLEC10A Antibody - N-terminal region - References**

Suzuki N., et al. J. Immunol. 156:128-135(1996).  
Ota T., et al. Nat. Genet. 36:40-45(2004).