

PEX11G Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15526**Specification**

PEX11G Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q96HA9
Other Accession	NM_080662 , NP_542393
Reactivity	Human, Rabbit
Predicted	Human, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27kDa KDa

PEX11G Antibody - C-terminal region - Additional Information**Gene ID** 92960**Other Names**

Peroxisomal membrane protein 11C, Peroxin-11C, Peroxisomal biogenesis factor 11C, Protein PEX11 homolog gamma, PEX11-gamma, PEX11G, PEX11C

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PEX11G antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

PEX11G Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PEX11G Antibody - C-terminal region - Protein Information**Name** PEX11G**Synonyms** PEX11C**Function**

Promotes membrane protrusion and elongation on the peroxisomal surface.

Cellular Location


Peroxisome membrane; Multi-pass membrane protein

PEX11G Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PEX11G Antibody - C-terminal region - Images



Host: Rabbit
Target Name: PEX11G
Sample Tissue: OVCAR-3 Whole cell lysate
s
Antibody Dilution: 1.0µg/ml

PEX11G Antibody - C-terminal region - References

Tanaka A.,et al.Biochem. Biophys. Res. Commun. 300:819-823(2003).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Li X.,et al.Mol. Cell. Biol. 22:8226-8240(2002).