

**MEX3A Antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI15713****Specification**

---

**MEX3A Antibody - N-terminal region - Product Information**

|                   |                                                                |
|-------------------|----------------------------------------------------------------|
| Application       | WB                                                             |
| Primary Accession | <a href="#">A1L020</a>                                         |
| Other Accession   | <a href="#">NM_001093725</a> , <a href="#">NP_001087194</a>    |
| Reactivity        | Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog |
| Predicted         | Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog |
| Host              | Rabbit                                                         |
| Clonality         | Polyclonal                                                     |
| Calculated MW     | 54kDa kDa                                                      |

**MEX3A Antibody - N-terminal region - Additional Information****Gene ID** 92312**Alias Symbol** MEX-3A, RKHD4**Other Names**

RNA-binding protein MEX3A, RING finger and KH domain-containing protein 4, MEX3A, RKHD4

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-MEX3A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

MEX3A Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**MEX3A Antibody - N-terminal region - Protein Information****Name** MEX3A**Synonyms** RKHD4**Function**

RNA binding protein, may be involved in post-transcriptional regulatory mechanisms.

**Cellular Location**

Cytoplasm. Nucleus. Cytoplasm, P-body. Note=Predominantly expressed in the cytoplasm and shuttles between the cytoplasm and the nucleus through the CRM1 export pathway


**Tissue Location**

Highest levels found in fetal brain and testis. Detected also in thymus, salivary gland and uterus

**MEX3A Antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**MEX3A Antibody - N-terminal region - Images**

Host: Rabbit  
Target Name: MEX3A  
Sample Tissue: Fetal Brain lysates  
Antibody Dilution: 1.0µg/ml

**MEX3A Antibody - N-terminal region - References**

Buchet-Poyau K., et al. Nucleic Acids Res. 35:1289-1300(2007).