

ARHGAP33 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15721**Specification**

ARHGAP33 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	K7EMC2
Reactivity	Human, Mouse, Rat, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Pig, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	92kDa KDa

ARHGAP33 Antibody - C-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ARHGAP33 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


ARHGAP33 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ARHGAP33 Antibody - C-terminal region - Protein Information**ARHGAP33 Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ARHGAP33 Antibody - C-terminal region - Images



168 kDa
144 kDa
90 kDa
65 kDa
40 kDa

Host: Rabbit

Target Name: ARHGAP33

Sample Tissue: 721_B Whole cell lysate

s

Antibody Dilution: 1.0µg/ml

ARHGAP33 Antibody - C-terminal region - Citations

- [Reduced mastication during growth inhibits cognitive function by affecting trigeminal ganglia and modulating Wnt signaling pathway and ARHGAP33 molecular transmission](#)