

CCDC13 Antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15742**Specification**

CCDC13 Antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8IYE1
Other Accession	NM_144719 , NP_653320
Reactivity	Human, Horse, Bovine
Predicted	Human, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	80kDa KDa

CCDC13 Antibody - N-terminal region - Additional Information**Gene ID** 152206**Other Names**

Coiled-coil domain-containing protein 13, CCDC13 (HGNC:26358)

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CCDC13 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CCDC13 Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CCDC13 Antibody - N-terminal region - Protein Information**Name** CCDC13 ([HGNC:26358](#))**Function**

Required for primary cilia formation and promotes the localization of the ciliopathy protein BBS4 to both centriolar satellites and cilia.

Cellular Location


Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite
Cytoplasm, cytoskeleton, cilium basal body

CCDC13 Antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCDC13 Antibody - N-terminal region - Images



Host: Rabbit
Target Name: CCDC13
Sample Tissue: Fetal Liver lysates
Antibody Dilution: 1.0µg/ml

CCDC13 Antibody - N-terminal region - References

Muzny D.M., et al. Nature 440:1194-1198(2006).
Staples C.J., et al. J. Cell Sci. 127:2910-2919(2014).