

OR4M2 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15772**Specification**

OR4M2 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8NGB6
Other Accession	NP_001004719
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34kDa kDa

OR4M2 Antibody - C-terminal region - Additional Information**Gene ID** 390538**Alias Symbol** OR4M2,
Other Names
Olfactory receptor 4M2, Olfactory receptor OR15-3, OR4M2**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 µl of distilled water. Final Anti-OR4M2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

OR4M2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

OR4M2 Antibody - C-terminal region - Protein Information**Name** OR4M2**Function**

Odorant receptor.

Cellular Location


Cell membrane; Multi-pass membrane protein.

OR4M2 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR4M2 Antibody - C-terminal region - Images



Host: Rabbit
Target Name: OR4M2
Sample Tissue: 786-0 Whole cell lysate
Antibody Dilution: 1.0µg/ml

OR4M2 Antibody - C-terminal region - Background

Odorant receptor.

OR4M2 Antibody - C-terminal region - References

Suwa M.,et al.Submitted (JUL-2001) to the EMBL/GenBank/DDBJ databases.
Zody M.C.,et al.Nature 440:671-675(2006).
Malnic B.,et al.Proc. Natl. Acad. Sci. U.S.A. 101:2584-2589(2004).
Malnic B.,et al.Proc. Natl. Acad. Sci. U.S.A. 101:7205-7205(2004).