

NUDT19 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15773**Specification**

NUDT19 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	A8MXV4
Other Accession	NP_001099040
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa

NUDT19 Antibody - C-terminal region - Additional Information**Gene ID** 390916**Alias Symbol** **NUDT19,**
Other Names

Nucleoside diphosphate-linked moiety X motif 19, mitochondrial, Nudix motif 19, 3.6.1.-, NUDT19

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 µl of distilled water. Final Anti-NUDT19 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

NUDT19 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NUDT19 Antibody - C-terminal region - Protein Information**Name** NUDT19**Function**

Fatty acyl-coenzyme A (CoA) diphosphatase that hydrolyzes fatty acyl-CoA to yield acyl-4'-phosphopantetheine and adenosine 3',5'- biphosphate (By similarity). Mediates the hydrolysis of a wide range of CoA esters, including choloyl-CoA and branched-chain fatty-acyl-CoA esters and at low substrate concentrations medium and long-chain fatty-acyl-CoA esters are the primary substrates (By similarity). Highest activity seen with medium-chain acyl-CoA esters and higher rates of activity seen with the unsaturated acyl-CoA esters compared with the saturated esters (By similarity). Exhibits decapping activity towards dpCoA-capped RNAs in vitro (By similarity).


Cellular Location

Peroxisome {ECO:0000250|UniProtKB:P11930}.

NUDT19 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NUDT19 Antibody - C-terminal region - Images

Host: Rabbit
Target Name: NUDT19
Sample Tissue: RPMI-8226 Whole cell lysate
s
Antibody Dilution: 1.0µg/ml

NUDT19 Antibody - C-terminal region - Background

Coenzyme A diphosphatase that mediates the hydrolysis of a wide range of CoA esters, including choloyl-CoA and branched- chain fatty-acyl-CoA esters. At low substrate concentrations medium and long-chain fatty-acyl-CoA esters are the primary substrates (By similarity).

NUDT19 Antibody - C-terminal region - References

Grimwood J.,et al.Nature 428:529-535(2004).
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).