

USP17L3 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15993**Specification**

USP17L3 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	A6NCW0
Other Accession	NM_001256871 , NP_001243800
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59kDa KDa

USP17L3 Antibody - C-terminal region - Additional Information**Gene ID** 645836**Alias Symbol** **USP17B, USP17F****Other Names**

Ubiquitin carboxyl-terminal hydrolase 17-like protein 3, 3.4.19.12, Deubiquitinating enzyme 17-like protein 3, Ubiquitin thioesterase 17-like protein 3, Ubiquitin-specific-processing protease 17-like protein 3, USP17L3, USP17B, USP17F

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 µl of distilled water. Final Anti- antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

USP17L3 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

USP17L3 Antibody - C-terminal region - Protein Information**Name** USP17L3**Synonyms** USP17B, USP17F**Function**

Deubiquitinating enzyme that removes conjugated ubiquitin from specific proteins to regulate different cellular processes that may include cell proliferation, progression through the cell cycle, apoptosis, cell migration, and the cellular response to viral infection.

Cellular Location


Nucleus. Endoplasmic reticulum

USP17L3 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

USP17L3 Antibody - C-terminal region - Images



Host: Rabbit

Target Name: USP17L3

Sample Tissue: Jurkat Whole cell lysate

s

Antibody Dilution: 1.0µg/ml

USP17L3 Antibody - C-terminal region - Background

Deubiquitinating enzyme that removes conjugated ubiquitin from specific proteins to regulate different cellular processes that may include cell proliferation, progression through the cell cycle, apoptosis, cell migration, and the cellular response to viral infection.

USP17L3 Antibody - C-terminal region - References

Nusbaum C.,et al.Nature 439:331-335(2006).
Burrows J.F.,et al.Genomics 85:524-529(2005).
Shin J.-M.,et al.BMC Genomics 7:292-292(2006).