10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HOXC13 Antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI16059

Specification

HOXC13 Antibody - middle region - Product Information

Application	**WB**
Primary Accession	P31276
Other Accession	NM_017410, NP_059106
Reactivity	**Human, Mouse, Rat, Rabbit, Pig, Sheep, Horse, Bovine, Neisseria Gonorrhoeae, Guinea Pig, Dog**
Predicted	**Human, Mouse, Rat, Rabbit, Pig, Sheep, Horse, Bovine, Neisseria Gonorrhoeae, Guinea Pig, Dog**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**36kDa KDa**

HOXC13 Antibody - middle region - Additional Information

Gene ID 3229

Alias Symbol	**HOX3, HOX3G**

Other Names
Homeobox protein Hox-C13, Homeobox protein Hox-3G, HOXC13, HOX3G

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 &mu, l of distilled water. Final Anti-HOXC13 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions
HOXC13 Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

HOXC13 Antibody - middle region - Protein Information

Name HOXC13

Synonyms HOX3G

Function
Transcription factor which plays a role in hair follicle differentiation. Regulates FOXQ1 expression and that of other hair- specific genes (By similarity).

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999


Cellular Location
Nucleus.

HOXC13 Antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

HOXC13 Antibody - middle region - Images



Host: Rabbit
Target Name: HOXC13
Sample Tissue: HepG2 Whole cell lysate
s
Antibody Dilution: 1.0µg/ml

HOXC13 Antibody - middle region - Background

Transcription factor which plays a role in hair follicle differentiation. Regulates FOXQ1 expression and that of other hair-specific genes (By similarity).

HOXC13 Antibody - middle region - References

de Stanchina E.,et al.J. Mol. Biol. 299:667-680(2000).
Li S.,et al.Submitted (JUN-2000) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (AUG-2003) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Acampora D.,et al.Nucleic Acids Res. 17:10385-10402(1989).