

STK24 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI16132**Specification**

STK24 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9Y6E0
Other Accession	NP_003567
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48kDa KDa

STK24 Antibody - C-terminal region - Additional Information**Gene ID** 8428**Alias Symbol** STK24, MST3, STK3,**Other Names**

Serine/threonine-protein kinase 24, 2.7.11.1, Mammalian STE20-like protein kinase 3, MST-3, STE20-like kinase MST3, Serine/threonine-protein kinase 24 36 kDa subunit, Mammalian STE20-like protein kinase 3 N-terminal, MST3/N, Serine/threonine-protein kinase 24 12 kDa subunit, Mammalian STE20-like protein kinase 3 C-terminal, MST3/C, STK24, MST3, STK3

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 µl of distilled water. Final Anti-STK24 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

STK24 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

STK24 Antibody - C-terminal region - Protein Information**Name** STK24 ([HGNC:11403](#))**Function**

Serine/threonine-protein kinase that acts on both serine and threonine residues and promotes apoptosis in response to stress stimuli and caspase activation. Mediates oxidative-stress-induced cell death by modulating phosphorylation of JNK1-JNK2 (MAPK8 and MAPK9), p38 (MAPK11, MAPK12, MAPK13 and MAPK14) during oxidative stress. Plays a role in a staurosporine-induced caspase-independent apoptotic pathway by regulating the nuclear translocation of AIFM1 and ENDOG and the DNase activity associated with ENDOG. Phosphorylates STK38L on 'Thr-442' and stimulates its kinase activity. In association with STK26 negatively regulates Golgi reorientation in

polarized cell migration upon RHO activation (PubMed:27807006). Also regulates cellular migration with alteration of PTPN12 activity and PXN phosphorylation: phosphorylates PTPN12 and inhibits its activity and may regulate PXN phosphorylation through PTPN12. May act as a key regulator of axon regeneration in the optic nerve and radial nerve. Part of the striatin-interacting phosphatase and kinase (STRIPAK) complexes. STRIPAK complexes have critical roles in protein (de)phosphorylation and are regulators of multiple signaling pathways including Hippo, MAPK, nuclear receptor and cytoskeleton remodeling. Different types of STRIPAK complexes are involved in a variety of biological processes such as cell growth, differentiation, apoptosis, metabolism and immune regulation (PubMed:18782753).

Cellular Location

Cytoplasm. Nucleus. Membrane. Note=The truncated form (MST3/N) translocates to the nucleus. Colocalizes with STK38L in the membrane

Tissue Location


Isoform A is ubiquitous. Isoform B is expressed in brain with high expression in hippocampus and cerebral cortex

STK24 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STK24 Antibody - C-terminal region - Images



Host: Rabbit
Target Name: STK24
Sample Tissue: 293T Whole Cell lysates
Antibody Dilution: 1.0µg/ml

STK24 Antibody - C-terminal region - Background

Serine/threonine-protein kinase that acts on both serine and threonine residues and promotes

apoptosis in response to stress stimuli and caspase activation. Mediates oxidative-stress- induced cell death by modulating phosphorylation of JNK1-JNK2 (MAPK8 and MAPK9), p38 (MAPK11, MAPK12, MAPK13 and MAPK14) during oxidative stress. Plays a role in a staurosporine-induced caspase-independent apoptotic pathway by regulating the nuclear translocation of AIFM1 and ENDOG and the DNase activity associated with ENDOG. Phosphorylates STK38L on 'Thr-442' and stimulates its kinase activity. Regulates cellular migration with alteration of PTPN12 activity and PXN phosphorylation: phosphorylates PTPN12 and inhibits its activity and may regulate PXN phosphorylation through PTPN12. May act as a key regulator of axon regeneration in the optic nerve and radial nerve.

STK24 Antibody - C-terminal region - References

Schinkmann K.,et al.J. Biol. Chem. 272:28695-28703(1997).
Zhou T.-H.,et al.J. Biol. Chem. 275:2513-2519(2000).
Dunham A.,et al.Nature 428:522-528(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Huang C.Y.,et al.J. Biol. Chem. 277:34367-34374(2002).