

PDCD8 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI16166**Specification**

PDCD8 antibody - N-terminal region - Product Information

Application	WB, IHC
Primary Accession	O95831-3
Other Accession	NM_145812 , NP_665811
Reactivity	Human, Sheep, Bovine, Dog
Predicted	Human, Sheep, Bovine, Dog, Cat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	66kDa KDa

PDCD8 antibody - N-terminal region - Additional Information

Alias Symbol	RP3-438D16.2, AIF, MGC111425, PDCD8, COXPD6
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PDCD8 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


PDCD8 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PDCD8 antibody - N-terminal region - Protein Information**PDCD8 antibody - N-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PDCD8 antibody - N-terminal region - Images



Human kidney



WB Suggested Anti-PDCD8 Antibody Titration: 0.2-1 μ g/ml

Positive Control: Jurkat cell lysate

AIFM1 is supported by BioGPS gene expression data to be expressed in Jurkat