

TNFRSF10C antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI16175**Specification**

TNFRSF10C antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O14798
Other Accession	NM_003841 , NP_003832
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25kDa KDa

TNFRSF10C antibody - N-terminal region - Additional Information**Gene ID** 8794

Alias Symbol CD263, DCR1, LIT, MGC149501, MGC149502, TRAILR3, TRID, TRAIL-R3, DCR1-TNFR

Other Names

Tumor necrosis factor receptor superfamily member 10C, Antagonist decoy receptor for TRAIL/Apo-2L, Decoy TRAIL receptor without death domain, Decoy receptor 1, DcR1, Lymphocyte inhibitor of TRAIL, TNF-related apoptosis-inducing ligand receptor 3, TRAIL receptor 3, TRAIL-R3, TRAIL receptor without an intracellular domain, CD263, TNFRSF10C, DCR1, LIT, TRAILR3, TRID

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-TNFRSF10C antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

TNFRSF10C antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TNFRSF10C antibody - N-terminal region - Protein Information**Name** TNFRSF10C**Synonyms** DCR1, LIT, TRAILR3, TRID**Function**

Receptor for the cytotoxic ligand TRAIL. Lacks a cytoplasmic death domain and hence is not capable of inducing apoptosis. May protect cells against TRAIL mediated apoptosis by competing

with TRAIL- R1 and R2 for binding to the ligand.

Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor.

Tissue Location


Higher expression in normal tissues than in tumor cell lines. Highly expressed in peripheral blood lymphocytes, spleen, skeletal muscle, placenta, lung and heart

TNFRSF10C antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TNFRSF10C antibody - N-terminal region - Images



WB Suggested Anti-TNFRSF10C Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: Human Stomach

TNFRSF10C antibody - N-terminal region - Background

Receptor for the cytotoxic ligand TRAIL. Lacks a cytoplasmic death domain and hence is not capable of inducing apoptosis. May protect cells against TRAIL mediated apoptosis by competing with TRAIL-R1 and R2 for binding to the ligand.

TNFRSF10C antibody - N-terminal region - References

MacFarlane M., et al. J. Biol. Chem. 272:25417-25420(1997).
Degli-Esposti M.A., et al. J. Exp. Med. 186:1165-1170(1997).
Pan G., et al. Science 277:815-818(1997).
Sheridan J.P., et al. Science 277:818-821(1997).
Schneider P., et al. FEBS Lett. 416:329-334(1997).