

RGS9 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI16193**Specification**

RGS9 antibody - N-terminal region - Product Information

Application	IHC, WB
Primary Accession	O75916
Other Accession	NM_003835 , NP_003826
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	49kDa KDa

RGS9 antibody - N-terminal region - Additional Information**Gene ID** 8787

Alias Symbol	PERRS, RGS9L
Other Names	
Regulator of G-protein signaling 9, RGS9, RGS9	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-RGS9 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

RGS9 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

RGS9 antibody - N-terminal region - Protein Information**Name** RGS9**Function**

Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to GNAT1. Involved in phototransduction; key element in the recovery phase of visual transduction (By similarity).

Cellular Location

[Isoform 3]: Membrane; Peripheral membrane protein. Note=Isoform 3 is targeted to the membrane via its interaction with RGS9BP.


Tissue Location

Highly expressed in the caudate and putamen, lower levels found in the hypothalamus and nucleus accumbens and very low levels in cerebellum. Not expressed in globus pallidus or cingulate cortex. Isoform 2 is expressed predominantly in pineal gland and retina. Isoform 3 is expressed in retina (abundant in photoreceptors)


RGS9 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RGS9 antibody - N-terminal region - Images

Human Liver



WB Suggested Anti-RGS9 Antibody Titration: 2.0µg/ml
Positive Control: HepG2 cell lysate

RGS9 antibody - N-terminal region - Background

Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby

driving them into their inactive GDP-bound form. Binds to G(t)-alpha. Involved in phototransduction; key element in the recovery phase of visual transduction (By similarity).

RGS9 antibody - N-terminal region - References

Granneman J.G.,et al.Mol. Pharmacol. 54:687-694(1998).

Zhang K.,et al.Gene 240:23-34(1999).

Puhl H.L. III,et al.Submitted (MAR-2004) to the EMBL/GenBank/DDBJ databases.

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Zody M.C.,et al.Nature 440:1045-1049(2006).