

NR3C2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI16235**Specification**

NR3C2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	B0ZBF5
Other Accession	NM_001166104 , NP_001159576
Reactivity	Human, Mouse, Rat, Horse, Bovine
Predicted	Human, Mouse, Rat, Chicken, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	94kDa kDa

NR3C2 antibody - C-terminal region - Additional Information

Alias Symbol	MR, MCR, MLR, FLJ41052, NR3C2VIT, MGC133092, NR3C2
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NR3C2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


NR3C2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NR3C2 antibody - C-terminal region - Protein Information**NR3C2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NR3C2 antibody - C-terminal region - Images



WB Suggested Anti-NR3C2 Antibody Titration: 0.2-1 μ g/ml
ELISA Titer: 1:12500
Positive Control: PANC1 cell lysate