

NFIC antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI16239**Specification**

NFIC antibody - middle region - Product Information

Application	WB
Primary Accession	Q6FI30
Other Accession	NM_205843 , NP_995315
Reactivity	Human, Pig, Dog
Predicted	Human, Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55kDa KDa

NFIC antibody - middle region - Additional Information

Alias Symbol **CTF, CTF5, MGC20153, NF-I, NFI**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NFIC antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


NFIC antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

NFIC antibody - middle region - Protein Information**NFIC antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NFIC antibody - middle region - Images



90 kDa
65 kDa
40 kDa
31 kDa
22 kDa

WB Suggested Anti-NFIC Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: 721_B cell lysate

NFIC is supported by BioGPS gene expression data to be expressed in 721_B