

APLNR/ Apelin Receptor / APJ Antibody (Extracellular Domain)
Rabbit Polyclonal Antibody
Catalog # ALS10021**Specification**

APLNR/ Apelin Receptor / APJ Antibody (Extracellular Domain) - Product Information

Application	IHC-P
Primary Accession	P35414
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43kDa KDa
Dilution	IHC-P~~N/A

APLNR/ Apelin Receptor / APJ Antibody (Extracellular Domain) - Additional Information**Gene ID** 187**Other Names**

Apelin receptor, Angiotensin receptor-like 1, G-protein coupled receptor APJ, G-protein coupled receptor HG11, APLNR, AGTRL1, APJ

Target/Specificity

Human Apelin Receptor. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

APLNR/ Apelin Receptor / APJ Antibody (Extracellular Domain) is for research use only and not for use in diagnostic or therapeutic procedures.

APLNR/ Apelin Receptor / APJ Antibody (Extracellular Domain) - Protein Information**Name** APLNR ([HGNC:339](#))**Synonyms** AGTRL1, APJ**Function**G protein-coupled receptor for peptide hormones apelin (APLN) and apelin receptor early endogenous ligand (APELA/ELA), that plays a role in the regulation of normal cardiovascular function and fluid homeostasis (PubMed: [11090199](http://www.uniprot.org/citations/11090199), PubMed: [22810587](http://www.uniprot.org/citations/22810587), PubMed: [25639753](http://www.uniprot.org/citations/25639753), PubMed: [28137936](http://www.uniprot.org/citations/28137936), PubMed: [35817871](http://www.uniprot.org/citations/35817871), PubMed: [38428423](http://www.uniprot.org/citations/38428423))

target="_blank">38428423). When acting as apelin receptor, activates both G(i) protein pathway that inhibits adenylate cyclase activity, and the beta-arrestin pathway that promotes internalization of the receptor (PubMed:11090199, PubMed:25639753, PubMed:28137936, PubMed:35817871, PubMed:38428423). APLNR/APJ also functions as mechanoreceptor that is activated by pathological stimuli in a G-protein-independent fashion to induce beta-arrestin signaling, hence eliciting cardiac hypertrophy (PubMed:22810587, PubMed:38428423). However, the presence of apelin ligand blunts cardiac hypertrophic induction from APLNR/APJ on response to pathological stimuli (PubMed:22810587, PubMed:38428423). Plays a key role in early development such as gastrulation, blood vessels formation and heart morphogenesis by acting as a APELA receptor (By similarity). May promote angioblast migration toward the embryonic midline, i.e. the position of the future vessel formation, during vasculogenesis (By similarity). Promotes sinus venosus (SV)-derived endothelial cells migration into the developing heart to promote coronary blood vessel development (By similarity). Also plays a role in various processes in adults such as regulation of blood vessel formation, blood pressure, heart contractility and heart failure (PubMed:25639753, PubMed:28137936).

Cellular Location

Cell membrane. Note=After exposure to apelin (APLN), internalized from the cell surface into an endosomal recycling compartment, from where it is recycled to the cell membrane (By similarity). After exposure to apelin receptor early endogenous ligand (APELA), internalized from the cell surface into an endosomal recycling compartment, from where it is recycled to the cell membrane (PubMed:25639753). {ECO:0000250|UniProtKB:Q9JHG3, ECO:0000269|PubMed:25639753}

Tissue Location

Expressed in heart, brain, kidney, stomach, spleen, thymus, lung, ovary, small intestine and colon, adipose tissues and pancreas (PubMed:25639753, PubMed:8294032). Expressed in glial cells, astrocytes and neuronal subpopulations (PubMed:8294032). Expressed in embryonic (ESCs) and induced (iPSCs) pluripotent stem cells (PubMed:25639753).

Volume


50 µl

APLNR/ Apelin Receptor / APJ Antibody (Extracellular Domain) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


APLNR/ Apelin Receptor / APJ Antibody (Extracellular Domain) - Images



Human Brain, Cortex: Formalin-Fixed, Paraffin-Embedded (FFPE)



Anti-APLNR/ Apelin Receptor / APJ antibody IHC of human Brain, Glioblastoma.



Anti-APLNR/ Apelin Receptor / APJ antibody IHC of human Colon, Carcinoma.

APLNR/ Apelin Receptor / APJ Antibody (Extracellular Domain) - Background

Receptor for apelin coupled to G proteins that inhibit adenylate cyclase activity and plays a role in various processes in adults such as regulation of blood pressure, heart contractility, and heart failure. Also plays a key role in early development such as gastrulation and heart morphogenesis by acting as a receptor for APELA hormone. Alternative coreceptor with CD4 for HIV-1 infection; may be involved in the development of AIDS dementia.

APLNR/ Apelin Receptor / APJ Antibody (Extracellular Domain) - References

O'Dowd B.F., et al. Gene 136:355-360(1993).

Eggerickx D., et al. Submitted (JUN-1995) to the EMBL/GenBank/DDBJ databases.