

## **GPR161 Antibody (N-Terminus)**

Rabbit Polyclonal Antibody Catalog # ALS10089

### **Specification**

## **GPR161 Antibody (N-Terminus) - Product Information**

Application IHC-P, ICC
Primary Accession Q8N6U8
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 59kDa KDa
Dilution IHC-P~~N/A

### **GPR161 Antibody (N-Terminus) - Additional Information**

### **Gene ID 23432**

### **Other Names**

G-protein coupled receptor 161, G-protein coupled receptor RE2, GPR161

## **Target/Specificity**

Human GPR161. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

### **Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

#### **Precautions**

GPR161 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

# **GPR161 Antibody (N-Terminus) - Protein Information**

### Name GPR161

### **Function**

Key negative regulator of Shh signaling, which promotes the processing of GLI3 into GLI3R during neural tube development. Recruited by TULP3 and the IFT-A complex to primary cilia and acts as a regulator of the PKA-dependent basal repression machinery in Shh signaling by increasing cAMP levels, leading to promote the PKA-dependent processing of GLI3 into GLI3R and repress the Shh signaling. In presence of SHH, it is removed from primary cilia and is internalized into recycling endosomes, preventing its activity and allowing activation of the Shh signaling. Its ligand is unknown (By similarity).

### **Cellular Location**

Cell projection, cilium membrane; Multi-pass membrane protein. Cell membrane; Multi-pass



membrane protein. Note=Mainly localizes to primary cilium in a TULP3 and IFT-A complex-dependent manner. In presence of SHH, it is removed from primary cilia and is internalized into recycling endosomes and is apparently not degraded (By similarity).


**Volume** 50 μl

# **GPR161 Antibody (N-Terminus) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# GPR161 Antibody (N-Terminus) - Images



Anti-GPR161 antibody IHC of human Lung, Adenocarcinoma.





Anti-GPR161 antibody ALS10089 IHC of human brain, pituitary.

## GPR161 Antibody (N-Terminus) - Background

Key negative regulator of Shh signaling, which promotes the processing of GLI3 into GLI3R during neural tube development. Recruited by TULP3 and the IFT-A complex to primary cilia and acts as a regulator of the PKA-dependent basal repression machinery in Shh signaling by increasing cAMP levels, leading to promote the PKA-dependent processing of GLI3 into GLI3R and repress the Shh signaling. In presence of SHH, it is removed from primary cilia and is internalized into recycling endosomes, preventing its activity and allowing activation of the Shh signaling. Its ligand is unknown (By similarity).

# **GPR161 Antibody (N-Terminus) - References**

Warren C.N.,et al.Submitted (APR-2003) to the EMBL/GenBank/DDBJ databases. Raming K.,et al.Recept. Channels 6:141-151(1998). Ota T.,et al.Nat. Genet. 36:40-45(2004). Gregory S.G.,et al.Nature 441:315-321(2006).