

**NMBR Antibody (C-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS10177****Specification**

---

**NMBR Antibody (C-Terminus) - Product Information**

Application	IHC-P
Primary Accession	<a href="#">P28336</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43kDa KDa
Dilution	IHC-P~~N/A

**NMBR Antibody (C-Terminus) - Additional Information****Gene ID** 4829**Other Names**

Neuromedin-B receptor, NMB-R, Epididymis tissue protein Li 185a, Neuromedin-B-preferring bombesin receptor, NMBR

**Target/Specificity**

Human NMBR. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

**Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

**Precautions**

NMBR Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**NMBR Antibody (C-Terminus) - Protein Information****Name** NMBR**Function**

Receptor for neuromedin-B (PubMed:<a href="http://www.uniprot.org/citations/1655761" target="\_blank">1655761</a>). Contributes to the maintenance of basal sigh rate through signaling in the pre- Botzinger complex, a cluster of several thousand neurons in the ventrolateral medulla responsible for inspiration during respiratory activity (By similarity). Contributes to the induction of sneezing following exposure to chemical irritants or allergens which causes release of NMB by nasal sensory neurons and activation of NMBR- expressing neurons in the sneeze-evoking region of the brainstem (By similarity). These in turn activate neurons of the caudal ventral respiratory group, giving rise to the sneezing response (By similarity). Contributes to induction of acute itch, possibly through its activation on dorsal root ganglion neurons by the NMB peptide (By similarity). Plays a role in the innate immune response to influenza A virus infection by enhancing

interferon alpha expression and reducing expression of IL6 (PubMed:<a href="http://www.uniprot.org/citations/31601264" target="\_blank">31601264</a>). Plays a role in CSF1-induced proliferation of osteoclast precursors by contributing to the positive regulation of the expression of the CSF1 receptor CSF1R (By similarity).

**Cellular Location**

Cell membrane; Multi-pass membrane protein

**Tissue Location**

Expressed in epididymis (at protein level).


**Volume**

50 µl


**NMBR Antibody (C-Terminus) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**NMBR Antibody (C-Terminus) - Images**

Anti-NMBR antibody IHC of human Ovary, Carcinoma.



Anti-NMBR antibody IHC of human Skin, Melanoma.

#### **NMBR Antibody (C-Terminus) - Background**

Receptor for neuromedin-B.

#### **NMBR Antibody (C-Terminus) - References**

Corjay M.H.,et al.J. Biol. Chem. 266:18771-18779(1991).

Li J.,et al.Mol. Cell. Proteomics 9:2517-2528(2010).

Mungall A.J.,et al.Nature 425:805-811(2003).

Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.