

TAAR6 Antibody (Cytoplasmic Domain)

Rabbit Polyclonal Antibody Catalog # ALS10540

Specification

TAAR6 Antibody (Cytoplasmic Domain) - Product Information

Application IHC
Primary Accession O96RI8
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 38kDa KDa

TAAR6 Antibody (Cytoplasmic Domain) - Additional Information

Gene ID 319100

Other Names

Trace amine-associated receptor 6, TaR-6, Trace amine receptor 6, Trace amine receptor 4, TaR-4, TAAR6, TA4, TAR4, TRAR4

Target/Specificity

Human TAAR6. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

TAAR6 Antibody (Cytoplasmic Domain) is for research use only and not for use in diagnostic or therapeutic procedures.

TAAR6 Antibody (Cytoplasmic Domain) - Protein Information

Name TAAR6

Synonyms TA4, TAR4, TRAR4

Function

Orphan receptor. Could be a receptor for trace amines. Trace amines are biogenic amines present in very low levels in mammalian tissues. Although some trace amines have clearly defined roles as neurotransmitters in invertebrates, the extent to which they function as true neurotransmitters in vertebrates has remained speculative. Trace amines are likely to be involved in a variety of physiological functions that have yet to be fully understood.

Cellular Location

Cell membrane; Multi-pass membrane protein.



Tissue Location

Expressed at low abundance in various brain tissues, as well as in fetal liver, but not in the cerebellum or placenta. In the brain, comparable levels of expression in basal ganglia, frontal cortex, substantia nigra, amygdala and hippocampus, highest expression in hippocampus and lowest expression in basal ganglia.


Volume 50 μl

TAAR6 Antibody (Cytoplasmic Domain) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

TAAR6 Antibody (Cytoplasmic Domain) - Images




Anti-TAAR6 antibody IHC of human Brain, Glioblastoma.




Anti-TAAR6 antibody ALS10540 IHC of human kidney, renal tubules.





Anti-TAAR6 antibody IHC of human Ovary, Carcinoma.



Anti-TAAR6 antibody IHC of human Skin, Melanoma.

TAAR6 Antibody (Cytoplasmic Domain) - Background

Orphan receptor. Could be a receptor for trace amines. Trace amines are biogenic amines present in very low levels in mammalian tissues. Although some trace amines have clearly defined roles as neurotransmitters in invertebrates, the extent to which they function as true neurotransmitters in vertebrates has remained speculative. Trace amines are likely to be involved in a variety of physiological functions that have yet to be fully understood.

TAAR6 Antibody (Cytoplasmic Domain) - References

Borowsky B.,et al.Proc. Natl. Acad. Sci. U.S.A. 98:8966-8971(2001). Kopatz S.A.,et al.Submitted (NOV-2002) to the EMBL/GenBank/DDBJ databases. Mungall A.J.,et al.Nature 425:805-811(2003). Duan J.,et al.Am. J. Hum. Genet. 75:624-638(2004).