

NPFF2 / NPFFR2 Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS10643**Specification**

NPFF2 / NPFFR2 Antibody (C-Terminus) - Product Information

Application	IHC-P
Primary Accession	Q9Y5X5
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60kDa KDa
Dilution	IHC-P~~N/A

NPFF2 / NPFFR2 Antibody (C-Terminus) - Additional Information**Gene ID** 10886**Other Names**

Neuropeptide FF receptor 2, G-protein coupled receptor 74, G-protein coupled receptor HLWAR77, Neuropeptide G-protein coupled receptor, NPFFR2, GPR74, NPFF2, NPGPR

Target/Specificity

Human NPFFR2. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

NPFF2 / NPFFR2 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

NPFF2 / NPFFR2 Antibody (C-Terminus) - Protein Information**Name** NPFFR2 ([HGNC:4525](#))**Synonyms** GPR74, NPFF2, NPGPR**Function**

Receptor for NPAF (A-18-F-amide) and NPFF (F-8-F-amide) neuropeptides, also known as morphine-modulating peptides. Can also be activated by a variety of naturally occurring or synthetic FMRF-amide like ligands. This receptor mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system.

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location

Isoform 1 is abundant in placenta. Relatively highly expressed in thymus, testis, and small intestine. Expressed at low levels in several tissues including spleen, prostate, brain, heart, ovary, colon, kidney, lung, liver and pancreas and not expressed in skeletal muscle and leukocytes. Isoform 2 expression is highest in placenta (but at relatively low level compared to isoform 1). Very low level of expression in numerous tissues including adipose tissue and many brain regions. Isoform 3 is expressed in brain and heart and, at lower levels, in kidney, liver, lung and pancreas


Volume

50 μ l


NPFF2 / NPFFR2 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NPFF2 / NPFFR2 Antibody (C-Terminus) - Images

Anti-NPFF2 / NPFFR2 antibody IHC of human testis.



Anti-NPFF2 / NPFFR2 antibody IHC of human colon.

NPFF2 / NPFFR2 Antibody (C-Terminus) - Background

Receptor for NPAF (A-18-F-amide) and NPFF (F-8-F-amide) neuropeptides, also known as morphine-modulating peptides. Can also be activated by a variety of naturally occurring or synthetic FMRF-amide like ligands. This receptor mediates its action by association with G proteins that activate a phosphatidylinositol- calcium second messenger system.

NPFF2 / NPFFR2 Antibody (C-Terminus) - References

Cikos S.,et al.Biochem. Biophys. Res. Commun. 256:352-356(1999).
Elshourbagy N.A.,et al.J. Biol. Chem. 275:25965-25971(2000).
Bonini J.A.,et al.J. Biol. Chem. 275:39324-39331(2000).
Parker R.M.C.,et al.Brain Res. Mol. Brain Res. 77:199-208(2000).
Liu Q.,et al.Submitted (DEC-2000) to the EMBL/GenBank/DDBJ databases.