

GPCRW / GPR18 Antibody (Extracellular Domain)
Rabbit Polyclonal Antibody
Catalog # ALS10718**Specification**

GPCRW / GPR18 Antibody (Extracellular Domain) - Product Information

Application	IHC-P
Primary Accession	Q14330
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38kDa KDa
Dilution	IHC-P~~N/A

GPCRW / GPR18 Antibody (Extracellular Domain) - Additional Information**Gene ID** 2841**Other Names**

N-arachidonyl glycine receptor, NAGly receptor, G-protein coupled receptor 18, GPR18, GPCRW

Target/Specificity

Human GPR18. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except GPR174 (44%).

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

GPCRW / GPR18 Antibody (Extracellular Domain) is for research use only and not for use in diagnostic or therapeutic procedures.

GPCRW / GPR18 Antibody (Extracellular Domain) - Protein Information**Name** GPR18**Synonyms** GPCRW**Function**

G protein-coupled receptor (GPCR) that plays a role in diverse physiological processes particularly within the immune and nervous systems (PubMed:21732409, PubMed:26195725). Becomes active when triggered by various endogenous ligands including endocannabinoid N- arachidonyl glycine (NAGly), delta-9-tetrahydrocannabinol or resolvin D2/RvD2 derived from the omega-3 fatty acid docosahexaenoic acid (DHA) (PubMed:16844083, PubMed:24762058, PubMed:26195725).

target="_blank">26195725, PubMed:27572937). Upon RvD2 binding, facilitates the resolution of inflammation, aiding in tissue repair and homeostasis. Mechanistically, RvD2 ligation initiates Gα_i protein coupling, activation of cAMP-PKA signaling pathway and phosphorylation of STAT3, leading to RvD2-stimulated macrophage phagocytosis (PubMed:27994074). Mediates NAGly-induced process of reorganization of actin filaments and induction of acrosomal exocytosis (PubMed:27572937). Activation by N-arachidonoyl glycine (NAGly) can also induce apoptosis in macrophages (By similarity). Plays a role in homeostasis of CD8⁺ subsets of intraepithelial lymphocytes (IELs) (CD8αα and CD8αβ IELs) in small intestine by supporting preferential migration of CD8αα T-cells to intraepithelial compartment over lamina propria compartment, and by mediating their reconstitution into small intestine after bone marrow transplant (By similarity). Also participates in hypotensive responses, mediating reduction in intraocular and blood pressure (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane

Tissue Location

Expressed in midpiece of spermatozoon (at protein level) (PubMed:27572937). Most abundant in testis and spleen (PubMed:16844083). Highly expressed in CD4 and CD8-positive T-cells as well as CD19-positive B-cells (PubMed:16844083)

Volume


50 µl

GPCRW / GPR18 Antibody (Extracellular Domain) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GPCRW / GPR18 Antibody (Extracellular Domain) - Images



Anti-GPR18 antibody ALS10718 IHC of human testis.

GPCRW / GPR18 Antibody (Extracellular Domain) - Background

Receptor for N-arachidonyl glycine. The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase. May contribute to regulation of the immune system.

GPCRW / GPR18 Antibody (Extracellular Domain) - References

- Gantz I.,et al.Genomics 42:462-466(1997).
Kohno M.,et al.Biochem. Biophys. Res. Commun. 347:827-832(2006).
Xu X.,et al.Submitted (MAY-2000) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (AUG-2003) to the EMBL/GenBank/DDBJ databases.
Dunham A.,et al.Nature 428:522-528(2004).