

**FZD6 / Frizzled 6 Antibody (N-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS10772****Specification**

---

**FZD6 / Frizzled 6 Antibody (N-Terminus) - Product Information**

Application	IHC, ICC
Primary Accession	<a href="#">O60353</a>
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	79kDa KDa

**FZD6 / Frizzled 6 Antibody (N-Terminus) - Additional Information****Gene ID** 8323**Other Names**

Frizzled-6, Fz-6, hFz6, FZD6

**Target/Specificity**

Human Frizzled-6. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except FZD3 (50%).

**Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

**Precautions**

FZD6 / Frizzled 6 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**FZD6 / Frizzled 6 Antibody (N-Terminus) - Protein Information****Name** FZD6**Function**

Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues. Together with FZD3, is involved in the neural tube closure and plays a role in the regulation of the establishment of planar cell polarity (PCP), particularly in the orientation of asymmetric bundles of stereocilia on the apical faces of a subset of auditory and vestibular sensory cells located in the inner ear (By similarity).

**Cellular Location**

Membrane {ECO:0000250|UniProtKB:Q61089}; Multi-pass membrane protein. Cell membrane {ECO:0000250|UniProtKB:Q61089}; Multi-pass membrane protein. Cell surface {ECO:0000250|UniProtKB:Q61089}. Apical cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q61089}; Multi-pass membrane protein. Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q61089}; Multi-pass membrane protein. Note=Colocalizes with FZD3 at the apical face of cells (By similarity). Localizes to the endoplasmic reticulum membrane in the presence of LMBR1L (By similarity). {ECO:0000250|UniProtKB:Q61089}

**Tissue Location**

Detected in adult heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, thymus, prostate, testis, ovary, small intestine and colon. In the fetus, expressed in brain, lung, liver and kidney


**Volume**

50 µl


**FZD6 / Frizzled 6 Antibody (N-Terminus) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**FZD6 / Frizzled 6 Antibody (N-Terminus) - Images**

Anti-Frizzled-6 antibody ALS10772 IHC of human pancreas.



Anti-Frizzled-6 antibody ALS10772 immunocytochemistry (ICC) staining of HEK293 human embryonic...

#### **FZD6 / Frizzled 6 Antibody (N-Terminus) - Background**

Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK- 3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues. Together with FZD3, is involved in the neural tube closure and plays a role in the regulation of the establishment of planar cell polarity (PCP), particularly in the orientation of asymmetric bundles of stereocilia on the apical faces of a subset of auditory and vestibular sensory cells located in the inner ear (By similarity).

#### **FZD6 / Frizzled 6 Antibody (N-Terminus) - References**

Tokuhara M.,et al.Biochem. Biophys. Res. Commun. 243:622-627(1998).  
Gazit A.,et al.Submitted (JUN-1998) to the EMBL/GenBank/DDBJ databases.  
Tanner S.M.,et al.Proc. Natl. Acad. Sci. U.S.A. 98:13901-13906(2001).  
Suwa M.,et al.Submitted (JUL-2001) to the EMBL/GenBank/DDBJ databases.  
Ota T.,et al.Nat. Genet. 36:40-45(2004).