

Ceramide Kinase / CERK Antibody (Internal)
Rabbit Polyclonal Antibody
Catalog # ALS10774**Specification**

Ceramide Kinase / CERK Antibody (Internal) - Product Information

Application	IHC-P
Primary Accession	Q8TCT0
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60kDa KDa
Dilution	IHC-P~~N/A

Ceramide Kinase / CERK Antibody (Internal) - Additional Information**Gene ID** 64781**Other Names**

Ceramide kinase, hCERK, 2.7.1.138, Acylsphingosine kinase, Lipid kinase 4, LK4, CERK, KIAA1646

Target/Specificity

Human CERK. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

Ceramide Kinase / CERK Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

Ceramide Kinase / CERK Antibody (Internal) - Protein Information**Name** CERK**Synonyms** KIAA1646**Function**

Catalyzes specifically the phosphorylation of ceramide to form ceramide 1-phosphate (PubMed:11956206, PubMed:16269826, PubMed:19168031). Acts efficiently on natural and analog ceramides (C6, C8, C16 ceramides, and C8-dihydroceramide), to a lesser extent on C2- ceramide and C6-dihydroceramide, but not on other lipids, such as various sphingosines (PubMed:11956206, PubMed:16269826, PubMed:19168031).

target="_blank">19168031). Shows a greater preference for D-erythro isomer of ceramides (PubMed:16269826). Binds phosphoinositides (PubMed:19168031).

Cellular Location

Cytoplasm. Cell membrane; Peripheral membrane protein

Tissue Location

High level expression in heart, brain, skeletal muscle, kidney and liver; moderate in peripheral blood leukocytes and thymus; very low in spleen, small intestine, placenta and lung

Volume


50 μ l

Ceramide Kinase / CERK Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ceramide Kinase / CERK Antibody (Internal) - Images



Anti-CERK antibody ALS10774 IHC of human brain, cortex.

Ceramide Kinase / CERK Antibody (Internal) - Background

Catalyzes specifically the phosphorylation of ceramide to form ceramide 1-phosphate. Acts efficiently on natural and analog ceramides (C6, C8, C16 ceramides, and C8-dihydroceramide), to a lesser extent on C2-ceramide and C6-dihydroceramide, but not on other lipids, such as various sphingosines. Binds phosphoinositides.

Ceramide Kinase / CERK Antibody (Internal) - References

Sugiura M.,et al.J. Biol. Chem. 277:23294-23300(2002).
Van Veldhoven P.P.,et al.Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.
Collins J.E.,et al.Genome Biol. 5:R84.1-R84.11(2004).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Dunham I.,et al.Nature 402:489-495(1999).