

FZD9 / Frizzled 9 Antibody (N-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS10807**Specification**

FZD9 / Frizzled 9 Antibody (N-Terminus) - Product Information

Application	IHC-P
Primary Accession	O00144
Reactivity	Human, Rabbit, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	64kDa KDa
Dilution	IHC-P~~N/A

FZD9 / Frizzled 9 Antibody (N-Terminus) - Additional Information**Gene ID** 8326**Other Names**

Frizzled-9, Fz-9, hFz9, FzE6, CD349, FZD9, FZD3

Target/Specificity

Human FZD9 / Frizzled 9. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

FZD9 / Frizzled 9 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

FZD9 / Frizzled 9 Antibody (N-Terminus) - Protein Information**Name** FZD9**Synonyms** FZD3**Function**

Receptor for WNT2 that is coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes (By similarity). Plays a role in neuromuscular junction (NMJ) assembly by negatively regulating the clustering of acetylcholine receptors (AChR) through the beta-catenin canonical signaling pathway (By similarity). May play a role in neural progenitor cells (NPCs) viability through the beta-catenin canonical signaling pathway by negatively regulating cell cycle arrest leading to inhibition of neuron apoptotic process (PubMed: 27509850). During hippocampal development, regulates neuroblast proliferation and apoptotic cell death.

Controls bone formation through non canonical Wnt signaling mediated via ISG15. Positively regulates bone regeneration through non canonical Wnt signaling (By similarity).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q9R216}; Multi-pass membrane protein.
Note=Relocalizes DVL1 to the cell membrane leading to phosphorylation of DVL1 and AXIN1 relocalization to the cell membrane. {ECO:0000250|UniProtKB:Q8K4C8}

Tissue Location

Expressed predominantly in adult and fetal brain, testis, eye, skeletal muscle and kidney. Moderately expressed in pancreas, thyroid, adrenal cortex, small intestine and stomach Detected in fetal liver and kidney. Expressed in neural progenitor cells (PubMed:27509850).


Volume

50 µl


FZD9 / Frizzled 9 Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FZD9 / Frizzled 9 Antibody (N-Terminus) - Images

Anti-FZD9 / Frizzled 9 antibody IHC of human Brain, Glioblastoma.



Anti-FZD9 / Frizzled 9 antibody ALS10807 IHC of human brains, neurons and glia.



Anti-FZD9 / Frizzled 9 antibody IHC of human testis.

FZD9 / Frizzled 9 Antibody (N-Terminus) - Background

Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues.

FZD9 / Frizzled 9 Antibody (N-Terminus) - References

Wang Y.-K., et al. Hum. Mol. Genet. 6:465-472(1997).
Hillier L.W., et al. Nature 424:157-164(2003).
Tanaka S., et al. Proc. Natl. Acad. Sci. U.S.A. 95:10164-10169(1998).