

## **TAOK1 / TAO1 Antibody (C-Terminus)**

Rabbit Polyclonal Antibody Catalog # ALS10884

## **Specification**

## TAOK1 / TAO1 Antibody (C-Terminus) - Product Information

Application IHC-P, E
Primary Accession Q7L7X3

Reactivity Human, Mouse, Rabbit, Hamster, Monkey,

Pig, Chicken, Horse, Xenopus, Bovine, Dog

Host Rabbit
Clonality Polyclonal
Calculated MW 116kDa KDa
Dilution IHC-P~~N/A
E~~N/A

# TAOK1 / TAO1 Antibody (C-Terminus) - Additional Information

#### Gene ID 57551

#### **Other Names**

Serine/threonine-protein kinase TAO1, 2.7.11.1, Kinase from chicken homolog B, hKFC-B, MARK Kinase, MARKK, Prostate-derived sterile 20-like kinase 2, PSK-2, PSK2, Prostate-derived STE20-like kinase 2, Thousand and one amino acid protein kinase 1, TAOK1, hTAOK1, TAOK1, KIAA1361, MAP3K16, MARKK

## Target/Specificity

Human TAOK1. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

#### **Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

## **Precautions**

TAOK1 / TAO1 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

## TAOK1 / TAO1 Antibody (C-Terminus) - Protein Information

#### Name TAOK1

Synonyms KIAA1361, MAP3K16, MARKK

#### **Function**

Serine/threonine-protein kinase involved in various processes such as p38/MAPK14 stress-activated MAPK cascade, DNA damage response and regulation of cytoskeleton stability. Phosphorylates MAP2K3, MAP2K6 and MARK2. Acts as an activator of the p38/MAPK14 stress-activated MAPK cascade by mediating phosphorylation and subsequent activation of the



upstream MAP2K3 and MAP2K6 kinases. Involved in G-protein coupled receptor signaling to p38/MAPK14. In response to DNA damage, involved in the G2/M transition DNA damage checkpoint by activating the p38/MAPK14 stress-activated MAPK cascade, probably by mediating phosphorylation of MAP2K3 and MAP2K6. Acts as a regulator of cytoskeleton stability by phosphorylating 'Thr-208' of MARK2, leading to activate MARK2 kinase activity and subsequent phosphorylation and detachment of MAPT/TAU from microtubules. Also acts as a regulator of apoptosis: regulates apoptotic morphological changes, including cell contraction, membrane blebbing and apoptotic bodies formation via activation of the MAPK8/JNK cascade. Plays an essential role in the regulation of neuronal development in the central nervous system (PubMed:<a href="http://www.uniprot.org/citations/33565190" target="\_blank">33565190</a>/a>). Also plays a role in the regulation of neuronal migration to the cortical plate (By similarity).

# Cellular Location Cytoplasm.

#### **Tissue Location**


Highly expressed in the testis, and to a lower extent also expressed in brain, placenta, colon and skeletal muscle

# TAOK1 / TAO1 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## TAOK1 / TAO1 Antibody (C-Terminus) - Images



Anti-TAOK1 antibody ALS10884 IHC of human liver.

# TAOK1 / TAO1 Antibody (C-Terminus) - Background

Serine/threonine-protein kinase involved in various processes such as p38/MAPK14 stress-activated MAPK cascade, DNA damage response and regulation of cytoskeleton stability. Phosphorylates MAP2K3, MAP2K6 and MARK2. Acts as an activator of the p38/MAPK14





Tel: 858.875.1900 Fax: 858.875.1999

stress-activated MAPK cascade by mediating phosphorylation and subsequent activation of the upstream MAP2K3 and MAP2K6 kinases. Involved in G-protein coupled receptor signaling to p38/MAPK14. In response to DNA damage, involved in the G2/M transition DNA damage checkpoint by activating the p38/MAPK14 stress-activated MAPK cascade, probably by mediating phosphorylation of MAP2K3 and MAP2K6. Acts as a regulator of cytoskeleton stability by phosphorylating 'Thr-208' of MARK2, leading to activate MARK2 kinase activity and subsequent phosphorylation and detachment of MAPT/TAU from microtubules. Also acts as a regulator of apoptosis: regulates apoptotic morphological changes, including cell contraction, membrane blebbing and apoptotic bodies formation via activation of the MAPK8/INK cascade.

# TAOK1 / TAO1 Antibody (C-Terminus) - References

Yustein J.T., et al. Oncogene 22:6129-6141(2003). Jenkins S.G., et al. Submitted (JUL-2001) to the EMBL/GenBank/DDBJ databases. Matsuda A., et al. Patent number WO2004058805, 15-JUL-2004. Nagase T., et al. DNA Res. 7:65-73(2000). Ota T., et al. Nat. Genet. 36:40-45(2004).